arXiv:1006.3954v1 [math.KT] 20 Jun 2010

# Analytic formulas for topological degree of non-smooth mappings: the even-dimensional case

Magnus Goffeng

Department of Mathematical Sciences, Division of Mathematics
Chalmers university of Technology and University of Gothenburg

**Abstract**

Topological degrees of continuous mappings between manifolds of even dimension are studied in terms of index theory of pseudo-differential operators. The index formalism of non-commutative geometry is used to derive analytic integral formulas for the index of a 0:th order pseudo-differential operator twisted by a Hölder continuous vector bundle. The index formula gives an analytic formula for the degree of a Hölder continuous mapping between even-dimensional manifolds. The paper is an independent continuation of the paper *Analytic formulas for topological degree of non-smooth mappings: the odd-dimensional case.*

## Introduction

This paper is an independent continuation of the paper [8] where the degree of a mapping from the boundary of a strictly pseudo-convex domain was given in terms of an explicit integral formula involving the Szegö kernel. In this paper analytic formulas are given for general even-dimensional manifolds in terms of the signature operator. The classical approach to mapping degree is to define in an abstract way the degree of a continuous mapping between two compact connected oriented manifolds of the same dimension in terms of homology. If the function $f$ is differentiable, an analytic formula for the degree can be derived using Brouwer degree, see [12], or the more global picture of de Rham cohomology. Without differentiability conditions on $f$, the only known analytic degree formula beyond [8] is a formula of Connes which only holds in one dimension, see more in Chapter 2.$\alpha$ of [5]. Our aim is to find another formula for the degree, that is valid for a Hölder continuous function, by expressing the degree as the index of a pseudo-differential operator and using the approach of [5] and [8].

Throughout the paper we will use the idea that the Chern character extracts cohomological information of a continuous mapping $f : X \to Y$ between even dimensional manifolds from the induced mapping $f^* : K^0(Y) \to K^0(X)$. The $K$-theory is a topological invariant and the picture of the index mapping as a pairing in a local homology theory via Chern characters in Atiyah-Singers index theorem can be applied to more general classes of functions than the smooth functions. The homology theory present throughout all the index theory is the cyclic homology. For a Hölder continuous mapping $f : X \to Y$ of exponent $\alpha$

and an elliptic pseudo-differential operator $A$ of order at least $\alpha$, this idea can be read out from the commutativity of the diagram:

$$\begin{array}{ccccc} K_0(C^\infty(Y)) & \xrightarrow{f^*} & K_0(C^\alpha(X)) & \xrightarrow{\text{ind}_A} & \mathbb{Z} \\ \downarrow{\scriptstyle \text{ch}_Y} & & \downarrow{\scriptstyle \text{ch}_X} & & \downarrow \\ HC_{even}(C^\infty(Y)) & \xrightarrow{f^*} & HC_{even}(C^\alpha(X)) & \xrightarrow{\tilde{\mu}_A} & \mathbb{C} \end{array} \tag{1}$$

where the mapping $\tilde{\mu}_A : HC_{even}(C^\alpha(X)) \to \mathbb{C}$ is a cyclic cocycle on $C^\alpha(X)$ defined as the Connes-Chern character of the bounded $K$-homology class that $A$ defines, see more in [4] and [5]. The right-hand side of the diagram (1) is commutative by Connes' index formula, see Proposition 4 of Chapter IV.1 of [5]. The dimension in which the Chern character will take values depends on the Hölder exponent $\alpha$. More explicitly, for $2n$-dimensional manifolds, the cocycle $\tilde{\mu}_A$ can be chosen as a cyclic $2k$-cocycle for any $k > n/\alpha$.

To describe this idea more explicitely, when $E \to X$ is a smooth vector bundle defined by the smooth projection-valued function $p : X \to \mathcal{K}$, the index of the twisted pseudo-differential operator $A_E := p(A \otimes \text{id}_{\mathcal{K}})p$ can be calculated in terms of the de Rham cohomology using the Atiyah-Singer index formula as

$$\text{ind}\, A_E = \int_X \text{ch}[E] \wedge \text{ch}[A] \wedge Td(X),$$

where $Td(X)$ denotes the Todd class of the complexified tangent bundle. In particular, if $E \to Y$ is a line bundle on an even-dimensional manifold $Y$ such that $\text{ch}[E]$ only contains a constant term and a top-degree term and $f : X \to Y$ is smooth we can consider the line bundle $f^*E \to X$. Naturality of the Chern character implies the identity

$$\deg(f)\, \text{ch}_0[A] \int_Y \text{ch}_Y[E] = \text{ind}\, A_{f^*E} - \text{ind}\,(A).$$

In Theorem 2.2, we construct an explicit line bundle $E_Y \to Y$ satisfying the above conditions together with the condition $\int_Y \text{ch}_Y[E_Y] = 1$. In the correct analytic setting the above degree formula extends to Hölder continuous functions. The analytic setting we choose in Theorem 4.2 is to associate a Fredholm module $(\tilde{\pi}, \tilde{F}_A)$ with an elliptic pseudo-differential operator $A$ of order at least $\alpha$. The Fredholm module $(\tilde{\pi}, \tilde{F}_A)$ is $q$-summable over the algebra of Hölder continuous functions $C^\alpha(X)$, for any $q > \dim(X)/\alpha$. Thus the Connes-Chern character $\tilde{\mu}_A := \text{cc}_k(\tilde{\pi}, \tilde{F}_A)$ is well defined for dimensions $2k > \dim(X)/\alpha$. In Theorem 5.1 we take $A$ to be the signature operator and show that if $f : X \to Y$ is Hölder continuous the following analytic degree formula holds:

$$\deg(f) = \tilde{\mu}_A(\text{ch}_X[f^*E_Y]) - \text{sign}(X).$$

The drawback exhibited in [8], where the results were restricted to boundaries of strictly pseudo-convex domains in Stein manifolds, is not present in this paper. The restriction that $X$ and $Y$ must be even-dimensional does not really pose a problem since when $X$ and $Y$ are odd-dimensional we can consider the mapping $f \times \text{id} : X \times S^1 \to Y \times S^1$ instead which is a mapping between even-dimensional manifolds and $\deg(f) = \deg(f \times \text{id})$. The drawback of the degree formula in Theorem 5.1 is that it is in general quite hard to calculate explicit integral kernels for pseudo-differential operators.

# 1 $K$-theory and Connes' index formula

To formulate the calculation of mapping degrees in a setting fitting with non-smooth mappings, we need a framework for "differential geometry" where there are no classical differentials. The framework we will use is Alain Connes' non-commutative geometry, see [4] and [5]. We will recall some basic concepts of non-commutative geometry in this section.

The $K$-theory of a compact topological space $Y$ is defined as the Grothendieck group of the abelian semigroup of isomorphism classes of vector bundles under direct sum. The Serre-Swan theorem states a one-to-one correspondence between the isomorphism classes of vector bundles over a compact space $Y$ and projection valued functions $p : Y \to \mathcal{K}$, see [14]. Here $\mathcal{K}$ denotes the $C^*$-algebra of compact operators on some separable, infinite dimensional Hilbert space. Following the Serre-Swan theorem, an equivalent approach to $K$-theory is to use equivalence classes of projections $p \in C(Y) \otimes \mathcal{K}$. The $K$-theory is denoted by $K_0(C(Y))$. To read more about $K$-theory, see [2]. The formulation of $K$-theory in terms of projections can be defined for any algebra $\mathcal{A}$ as equivalence classes of projections $p \in \mathcal{A} \otimes \mathcal{K}$.

Clearly, the abelian group $K_0(\mathcal{A})$ depends covariantly on the algebra $\mathcal{A}$ so $K_0$ defines a functor. In particular, the functor $K_0$ has many properties making the $K$-theory of a $C^*$-algebra manageable to calculate, for instance; homotopy invariance, half exactness and stability under tensoring by a matrix algebra. Furthermore, a dense embedding of topological algebras $\mathcal{A}' \hookrightarrow \mathcal{A}$ which is isoradial induces an isomorphism on $K$-theory, see more in [6]. For instance, if $Y$ is a compact manifold all of the embeddings $C^\infty(Y) \subseteq C^\alpha(Y) \subseteq C(Y)$ induce isomorphisms on $K$-theory. Here $C^\alpha(Y)$ denotes the algebra of Hölder continuous functions of exponent $\alpha \in ]0, 1]$. The isomorphism $K_0(C(Y)) \cong K_0(C^\infty(Y))$ enables us to define the Chern character $\mathrm{ch} : K_0(C(Y)) \to H^{even}_{dR}(Y)$ by representing a class $[p] \in K_0(C(Y))$ by a smooth $p : Y \to \mathcal{K}$ and define

$$\mathrm{ch}[p] := \sum_{j=0}^{\infty} \frac{1}{(2\pi)^j j!} \mathrm{tr}(p\mathrm{d}p\mathrm{d}p)^j.$$

We choose the trace as the fiberwise operator trace in $\wedge^* T^*Y \otimes \mathcal{K}$ which is well defined since a compact projection is of finite rank. The term in the sum of degree $2j$ is denoted by $\mathrm{ch}_j[p]$.

However, we will need a Chern character defined on Hölder continuous projections. The homology theory fitting with index theory of more complicated geometries than smooth functions on smooth manifolds is cyclic homology. We will consider Connes' original definition of cyclic homology which simplifies the construction of the Chern character and the Chern-Connes character. We will let $\mathcal{A}$ denote a topological algebra and we will use the notation $\mathcal{A}^{\otimes k}$ for the $k$-th tensor power of $\mathcal{A}$. The Hochschild differential $b : \mathcal{A}^{\otimes k} \to \mathcal{A}^{\otimes k-1}$ is defined by

$$\begin{aligned} b(x_0 \otimes x_1 \otimes \cdots \otimes x_k \otimes x_{k+1}) &:= (-1)^{k+1} x_{k+1}x_0 \otimes x_1 \otimes \cdots \otimes x_k + \\ &+ \sum_{j=0}^{k} (-1)^j x_0 \otimes \cdots \otimes x_{j-1} \otimes x_j x_{j+1} \otimes x_{j+2} \otimes \cdots \otimes x_{k+1}. \end{aligned}$$

The cyclic permutation operator $\lambda : \mathcal{A}^{\otimes k} \to \mathcal{A}^{\otimes k}$ is defined as

$$\lambda(x_0 \otimes x_1 \otimes \cdot \otimes x_k) = (-1)^k x_k \otimes x_0 \otimes \cdots \otimes x_{k-1}.$$

We define a complex of $\mathbb{C}$-vector spaces $C_*^\lambda(\mathcal{A})$ by

$$C_k^\lambda(\mathcal{A}) := \mathcal{A}^{\otimes k+1}/(1-\lambda)\mathcal{A}^{\otimes k+1},$$

with differential given by $b$. The homology of the complex $C_*^\lambda(\mathcal{A})$ is called the cyclic homology of $\mathcal{A}$ and will be denoted by $HC_*(\mathcal{A})$. A cycle in $C_\lambda^k(\mathcal{A})$ will be called a cyclic $k$-cycle.

The complex $C_\lambda^k(\mathcal{A})$ is defined as the space of continuous linear functionals $\mu$ on $\mathcal{A}^{\otimes k+1}$ such that $\mu \circ \lambda = \mu$. The Hochschild coboundary operator $\mu \mapsto \mu \circ b$ makes $C_\lambda^*(\mathcal{A})$ into a complex. The cohomology of the complex $C_\lambda^*(\mathcal{A})$ will be denoted by $HC^*(\mathcal{A})$ and is called the cyclic cohomology of $\mathcal{A}$. Cyclic cohomology is an algebraic generalization of de Rham homology. The difference lies in that the dimension defines a grading on the de Rham theories, while the dimension defines a filtration on the cyclic theories. This difference can be explained by a Theorem of Connes [5] stating that if $X$ is a compact manifold, there is an isomorphism

$$HC^k(C^\infty(X)) \cong Z_k(X) \oplus \bigoplus_{j\geq 0} H^{dR}_{k-2j}(X), \tag{2}$$

where $Z_k(X)$ denotes the space of closed $k$-currents on $X$. The filtration on cyclic cohomology can be described by the linear mapping $S : HC^k(\mathcal{A}) \to HC^{k+2}(\mathcal{A})$ called the suspension operator or the periodicity operator. For a definition of the periodicity operator, see [5].

The Chern character $\mathrm{ch}_k : K_0(\mathcal{A}) \to HC_{2k}(\mathcal{A})$ in degree $2k$ is defined as in Proposition 3 of Chapter $III.3$ of [5] by

$$\mathrm{ch}_k[p] := (k!)^{-1}\mathrm{tr}\left(\underbrace{p \otimes p \otimes \cdots \otimes p \otimes p}_{2k+1 \quad \text{factors}}\right). \tag{3}$$

The additive pairing between $HC^{2k}(\mathcal{A})$ and the $K$-theory $K_0(\mathcal{A})$ is defined by

$$\langle \mu, x\rangle_k := \mu.\mathrm{ch}[x].$$

The choice of normalization implies that for a cohomology class in $HC^{2k}(\mathcal{A})$ represented by the cyclic cocycle $\mu$, the pairing satisfies

$$\langle S\mu, x\rangle_{k+1} = \langle \mu, x\rangle_k,$$

see Proposition 3 in Chapter III.3 of [5].

The homology theory dual to $K$-theory is $K$-homology. Analytic $K$-homology is described by Fredholm modules, a theory fitting well with index theory and cyclic cohomology. For $q \geq 1$, let $\mathcal{L}^q(\mathcal{H}) \subseteq \mathcal{B}(\mathcal{H})$ denote the ideal of Schatten class operators on a separable Hilbert space $\mathcal{H}$, so $T \in \mathcal{L}^q(\mathcal{H})$ if and only if $\mathrm{tr}((T^*T)^{q/2}) < \infty$. For $q \neq 2$ there is no exact description for an integral operator to belong to the Schatten class of order $q$. However, for $q > 2$ there exists a convenient sufficient condition on the kernel, found in [13]. We will return to this subject a little later.

A graded Hilbert space is a Hilbert space $\mathcal{H}$ equipped with an involutive mapping $\gamma$, that is, $\gamma^2 = 1$. While $\gamma$ is an involution, we can decompose

$\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-$, where $\mathcal{H}_\pm = \ker(\gamma \mp 1)$. An operator $T$ on $\mathcal{H}$ is called even if $T\mathcal{H}_\pm \subseteq \mathcal{H}_\pm$ and odd if $T\mathcal{H}_\pm \subseteq \mathcal{H}_\mp$. Suppose that $\mathcal{H}$ is a graded Hilbert space and $\pi : \mathcal{A} \to \mathcal{B}(\mathcal{H})$ is an even representation of a trivially graded $\mathbb{C}$-algebra $\mathcal{A}$. If $F \in \mathcal{B}(\mathcal{H})$ is an odd operator such that

$$F^2 = 1,\ F = F^* \quad \text{and} \quad [F, \pi(a)] \in \mathcal{L}^q(\mathcal{H})\ \forall a \in \mathcal{A}, \tag{4}$$

the pair $(\pi, F)$ is called a $q$-summable even Fredholm module. The conditions $F^2 = 1$ and $F = F^*$ simplifies many calculations, but in practice it is sufficient if they hold modulo $q$-summable operators. If we decompose the graded Hilbert space $\mathcal{H} = \mathcal{H}_+ \oplus \mathcal{H}_-$ into its even and odd part, the odd operator $F$ decomposes as:

$$F = \begin{pmatrix} 0 & F_+ \\ F_- & 0 \end{pmatrix}$$

where $F_+ : \mathcal{H}_- \to \mathcal{H}_+$ and $F_- : \mathcal{H}_+ \to \mathcal{H}_-$. Similarly we can decompose $\pi = \pi_+ \oplus \pi_-$ where $\pi_\pm : \mathcal{A} \to \mathcal{B}(\mathcal{H}_\pm)$ are representations. The first and second condition in (4) are equivalent to the conditions $F_+ = F_-^{-1} = F_-^*$ and the commutator condition is equivalent to

$$F_-\pi_+(a) - \pi_-(a)F_- \in \mathcal{L}^q(\mathcal{H}_+, \mathcal{H}_-) \quad \text{and} \quad F_+\pi_-(a) - \pi_+(a)F_+ \in \mathcal{L}^q(\mathcal{H}_-, \mathcal{H}_+).$$

If the pair $(\pi, F)$ satisfies the requirement in equation (4) but with $\mathcal{L}^q(\mathcal{H})$ replaced by $\mathcal{K}(\mathcal{H})$ the pair $(\pi, F)$ is a bounded even Fredholm module. The set of homotopy classes of bounded even Fredholm modules forms an abelian group under direct sum called the even analytic $K$-homology of $\mathcal{A}$ and is denoted by $K^0(\mathcal{A})$. For a more thorough presentation of Fredholm modules, e.g. Chapter VII and VIII of [2].

Following Definition 3 of Chapter IV.1 of [5] we define the Connes-Chern character $\mathrm{cc}_k(\pi, F)$ of a $q$-summable even Fredholm module $(\pi, F)$ for $2k \geq q$ as the cyclic $2k$-cocycle:

$$\mathrm{cc}_k(\pi, F)(a_0, a_1, \ldots, a_{2k}) := (-1)^k k!\, \mathrm{str}(\pi(a_0)[F, \pi(a_1)] \cdots [F, \pi(a_{2k})]),$$

where $\mathrm{str}(T) := \mathrm{tr}(\gamma T)$ for $T \in \mathcal{L}^1(\mathcal{H})$ and $\gamma$ denotes the grading on $\mathcal{H}$. This choice of normalization leads to $S\mathrm{cc}_k(\pi, F) = cc_{k+1}(\pi, F)$, see Proposition 2 of Chapter IV.1 of [5].

If $(\pi, F)$ is a bounded Fredholm module over $\mathcal{A}$ we can define the index mapping $\mathrm{ind}_F : K_0(\mathcal{A}) \to \mathbb{Z}$ as

$$\mathrm{ind}_F[p] := \mathrm{ind}\,((\pi_+ \otimes \mathrm{id})(p)F_+(\pi_- \otimes \mathrm{id})(p)),$$

where we represent $[p]$ by a finite-dimensional projection $p \in \mathcal{A} \otimes \mathcal{K}(\mathbb{C}^N)$ and we consider $(\pi_+ \otimes \mathrm{id})(p)F_+(\pi_- \otimes \mathrm{id})(p)$ as an operator

$$(\pi_+ \otimes \mathrm{id})(p)F_+(\pi_- \otimes \mathrm{id})(p) : (\pi_- \otimes \mathrm{id})(p)\left(\mathcal{H}_- \otimes \mathbb{C}^N\right) \to (\pi_+ \otimes \mathrm{id})(p)\left(\mathcal{H}_+ \otimes \mathbb{C}^N\right).$$

The association $[p] \times (\pi, F) \mapsto \mathrm{ind}_F(p)$ is homotopy invariant and defines a bilinear pairing $K_0(\mathcal{A}) \times K^0(\mathcal{A}) \to \mathbb{Z}$ which is non-degenerate after tensoring with $\mathbb{Q}$, see more in [2]. To simplify the notation, we suppress the dimension $N$ and identify $(\pi, F)$ with the Fredholm module $(\pi \otimes \mathrm{id}_{\mathcal{K}(\mathbb{C}^N)}, F \otimes \mathrm{id}_{\mathbb{C}^N})$.

**Theorem 1.1** (Proposition 4 of Chapter IV.1 of [5]). *If $(\pi, F)$ is a $q$-summable even Fredholm module and $2k \geq q$ the index mapping* $\mathrm{ind}_F$ *can be calculated as*

$$\mathrm{ind}_F[p] = \langle \mathrm{cc}_k(\pi, F), p\rangle_k.$$

In Theorem 1.1, the conditions (4) on the Fredholm module $(\pi, F)$ requires some caution. If for instance, we remove the condition $F^2 \neq 1$ one can choose $F$ such that $\mathrm{ind}\,(F_+) \neq 0$. On the other hand if we require $\pi$ to be unital, we have that $\mathrm{ind}\,(F_+) = \mathrm{ind}_F(1)$ but $\langle \mathrm{cc}_k(\pi, F), 1\rangle_k = 0$, therefore $\mathrm{ind}\,(F_+) \neq \langle \mathrm{cc}_k(\pi, F), 1\rangle_k$.

In the context of index theory, the periodicity operator $S$ plays the role of extending index formulas such as that in Theorem 1.1 to larger algebras. Suppose that $\mu$ is a cyclic $k$-cocycle on an algebra $\mathcal{A}_0$ which is a dense $*$-subalgebra of a $C^*$-algebra $\mathcal{A}$. The cyclic $k+2m$-cocycle $S^m\mu$ can sometimes be extended by continuity to a cyclic cocycle on a larger $*$-subalgebra $\mathcal{A}_0 \subseteq \mathcal{A}' \subseteq \mathcal{A}$. In [8] the properties of $S^m\mu$ were studied for $\Omega$ being a strictly pseudo-convex domain in a Stein manifold of complex dimension $n$ and $\mu$ being the cyclic $2n-1$-cocycle on $\mathcal{A}_0 = C^\infty(\partial\Omega)$ defined by

$$\mu := \sum_{k=0}^{n} S^k\omega_k,$$

where $\omega_k$ denotes the cyclic $2n-2k-1$-cocycle given by the Todd class $Td_k(\Omega)$ in degree $2k$ as

$$\omega_k(a_0, a_1, \ldots, a_{2n-2k-1}) := \int_{\partial\Omega} a_0 \mathrm{d}a_1 \wedge \mathrm{d}a_2 \wedge \cdots \wedge \mathrm{d}a_{2n-2k-1} \wedge Td_k(\Omega).$$

It was proved in [8] that the cyclic cocycle $S^m\mu$ defines the same cyclic cohomology class on $C^\infty(\partial\Omega)$ as

$$\tilde{\mu}(a_0, a_1, \ldots, a_{2n+2m-1}) := \tag{5}$$

$$:= \int_{\partial\Omega^{2n+2m-1}} \mathrm{tr}\left(a_0(z_0) \prod_{j=1}^{2n+2m-1} (a_j(z_j) - a_j(z_{j-1}))K_{\partial\Omega}(z_{j-1}, z_j)\right) \mathrm{d}V, \tag{6}$$

where $K_{\partial\Omega}$ denotes the Szegö kernel or the Henkin-Ramirez kernel. The cyclic cocycle $\tilde{\mu}$ is the *odd* Connes-Chern character of the Toeplitz operators on the Hardy space and $\tilde{\mu}$ extends to a cyclic cocycle on the algebra of Hölder continuous functions on $\partial\Omega$. We will later on use a cyclic cocycle of the form $\mu = \mathrm{cc}_k(\pi, F)$ and the periodicity operator to extend a formulation of the Atiyah-Singer index theorem to pseudo-differential operators twisted by a Hölder continuous vector bundle.

The index formula of Theorem 1.1 holds for $q$-summable Fredholm modules and to deal with the $q$-summability of pseudo-differential operators we will need the following theorem of Russo [13] to give a sufficient condition for an integral operator to be Schatten class. Following [3], when $X$ is a $\sigma$-finite measure space and $1 \leq p, q < \infty$, the mixed $(p, q)$-norm of a function $k : X \times X \to \mathbb{C}$ is defined by

$$\|k\|_{p,q} := \left(\int_X \left(\int_X |k(x, y)|^p \mathrm{d}x\right)^{\frac{q}{p}} \mathrm{d}y\right)^{\frac{1}{q}}.$$

The space of measurable functions with finite mixed $(p,q)$-norm is denoted by $L^{(p,q)}(X\times X)$. By Theorem 4.1 of [3] the space $L^{(p,q)}(X\times X)$ becomes a Banach space in the mixed $(p,q)$-norm which is reflexive if $1<p,q<\infty$. If a bounded operator $K$ has integral kernel $k$, the hermitian conjugate $K^*$ has integral kernel $k^*(x,y):=\overline{k(y,x)}$.

**Theorem 1.2** (Theorem 1 in [13])**.** *Suppose that $K:L^2(X)\to L^2(X)$ is a bounded operator given by an integral kernel $k$. If $2<q<\infty$*

$$\|K\|_{\mathcal{L}^q(L^2(X))}\leq(\|k\|_{q',q}\|k^*\|_{q',q})^{1/2}, \tag{7}$$

*where $q'=q/(q-1)$.*

In the statement of the Theorem in [13], the completely unnecessary assumption $k\in L^2(X\times X)$ is made. For the discussion on how to remove the condition $k\in L^2(X\times X)$ and the proof of the next Theorem, we refer to [8].

**Theorem 1.3.** *Suppose that $K_j:L^2(X)\to L^2(X)$ are operators with integral kernels $k_j$ for $j=1,\ldots,m$ such that $\|k_j\|_{q',q},\|k_j^*\|_{q',q}<\infty$ for certain $q>2$. Whenever $m\geq q$ the operator $K_1K_2\cdots K_m$ is a trace class operator and we have the trace formula*

$$\mathrm{tr}(K_1K_2\cdots K_m)=\int_{X^m}\left(\prod_{j=1}^m k_j(x_j,x_{j+1})\right)\mathrm{d}x_1\mathrm{d}x_2\cdots\mathrm{d}x_m,$$

*where we identify $x_{m+1}$ with $x_1$.*

# 2 The projection with Chern character being the volume form

In order to obtain a formulation of the degree as an index, we start by constructing a line bundle $E_Y$ over an arbitrary even dimensional manifold $Y$ such that the only non-constant term in $\mathrm{ch}[E_Y]$ is of top degree. The idea is to use the tautological line bundle over $S^2$ and the fact that $S^{2n}$ is the smashed products of $n$ copies of $S^2$ to define a line bundle over $S^{2n}$ for arbitrary $n$. Under the diffeomorphism $S^2\cong P^1_{\mathbb{C}}$, the projective complex line, we take the complex coordinate $z$ on $S^2$ corresponding to one of the affine charts. In the $z$-chart the tautological line bundle is defined by the projection valued function $p_0:\mathbb{C}\to M_2(\mathbb{C})$ which is given as

$$p_0(z):=\frac{1}{1+|z|^2}\begin{pmatrix}|z|^2 & z\\ \bar{z} & 1\end{pmatrix}.$$

If we define $v:\mathbb{C}\to\mathbb{C}^2$ by $v(z):=\frac{1}{\sqrt{1+|z|^2}}(z,1)$ then for $w\in\mathbb{C}^2$ we have that

$$p_0(z)w=\langle w,v(z)\rangle v(z). \tag{8}$$

It follows that the function $p_0$ satisfies $p_0^2=p_0^*=p_0$, so $p_0$ is a hermitian projection. Let us denote by $\omega$ the Fubini-Study metric on $S^2$, so

$$\omega:=\frac{i\mathrm{d}z\wedge\mathrm{d}\bar{z}}{(1+|z|^2)^2}.$$

A straightforward calulation shows that

$$\mathrm{ch}[p_0] = 1 + \frac{\omega}{2\pi}$$

In higher dimensions we define $p_T : \mathbb{C}^n \to M_{2^n}(\mathbb{C})$ by

$$p_T(z_1, z_2, \ldots, z_n) := p_0(z_1) \otimes p_0(z_2) \otimes \ldots \otimes p_0(z_n). \tag{9}$$

Let us use the notation

$$p_j(z_1, z_2, \ldots, z_n) := 1 \otimes \cdots \otimes 1 \otimes p_0(z_j) \otimes \ldots \otimes 1 \ldots \otimes 1,$$

with $p_0(z_j)$ in the $j$:th position. It follows that

$$\mathrm{ch}[p_T] = \mathrm{ch}[p_1 p_2 \cdots p_n] = \sum_{I \subseteq \{1,2,\ldots,n\}} (2\pi)^{-|I|} \bigwedge_{j \in I} \omega_j, \tag{10}$$

where $\omega_j$ denotes the Fubini-Study metric depending on the $j$:th variable $z_j$.

**Lemma 2.1.** *The projection $p_T$ extends to a projection valued $C^1$-function on $S^{2n}$ such that*

$$\mathrm{ch}[p_T] = 1 + \mathrm{d}V_{S^{2n}}$$

*in $H^{even}_{dR}(S^{2n})$.*

*Proof.* Since $H^{even}_{dR}(S^{2n})$ is spanned by the constant function and the volume form, it is sufficient to show that $\mathrm{ch}_0[p_T] = 1$ and $\mathrm{ch}_n[p_T] = \mathrm{d}V_{S^{2n}}$. The function $p_T$ takes values as a rank-one projection, therefore $\mathrm{ch}_0[p_T] = 1$. That $\mathrm{ch}_n[p_T] = \mathrm{d}V_{S^{2n}}$ follows from the following calculation

$$\int_{S^{2n}} \mathrm{ch}_k[p_T] = (2\pi)^{-n} \int_{\mathbb{C}^n} \bigwedge_{j=1}^{n} \omega_j =$$

$$= 2^n (2\pi)^{-n} \int_{\mathbb{C}^n} \prod_{j=1}^{n} \frac{\mathrm{d}V_{\mathbb{C}}(z_j)}{(1+|z_j|^2)^2} = \left( \int_{\mathbb{C}} \frac{2r\mathrm{d}r}{(1+r^2)^2} \right)^n = 1.$$

□

In the general case, let $Y$ be a compact connected orientable manifold of dimension $2n$. If we take an open subset $U$ of $Y$ with coordinates $(x_i)_{i=1}^{2n}$ such that

$$U = \{x : \sum_{i=1}^{2n} |x_i(x)|^2 < 1\},$$

the coordinates define a diffeomorphism $\nu : U \cong B_{2n}$. Let us also choose a diffeomorphism $\tau : B_{2n} \cong \mathbb{C}^n$. We can define the projection valued functions $p_Y : Y \to M_{2^n}(\mathbb{C})$ by

$$p_Y(x) := \begin{cases} p_T(\tau\nu(x)) & \text{for} \quad x \in U \\ p_T(\infty) & \text{for} \quad x \notin U \end{cases} \tag{11}$$

If we let $\tilde{\nu} : Y \to S^{2n}$ be the Lipschitz continuous function defined by

$$\tilde{\nu}(x) = \begin{cases} \tau(\nu(x)) & \text{for} \quad x \in U \\ \infty & \text{for} \quad x \notin U \end{cases} \tag{12}$$

the $C^1$-function $p_Y$ can be expressed as $p_Y = \tilde{\nu}^* p_T$.

**Theorem 2.2.** *If $Y$ is a compact connected orientable manifold of even dimension and $\mathrm{d}V_Y$ denotes the normalized volume form on $Y$, the projection $p_Y$ satisfies*

$$\mathrm{ch}[p_Y] = 1 + \mathrm{d}V_Y, \tag{13}$$

*in $H^{even}_{dR}(Y)$. Thus, if $f : X \to Y$ is a smooth mapping*

$$\deg(f) = \int_X f^* \mathrm{ch}[p_Y]$$

*Proof.* By Lemma 2.1 we have the identities

$$\int_Y \mathrm{ch}[p_Y] = \int_U \mathrm{ch}_n[p_Y] = \int_U \tilde{\nu}^* \mathrm{ch}_n[p_T] = \int_{S^{2n}} \mathrm{ch}_k[p_T] = 1.$$

Therefore we have the identity $\mathrm{ch}_n[p] = \mathrm{d}V_Y$. Since $\mathrm{ch}[p_Y] = 1 + \mathrm{ch}_n[p_Y]$ up to an exact form on $U$ and vanishes to first order at $\partial U$ the Theorem follows. □

Later on we will also need the Chern character of $p_Y$ in cyclic homology as is defined in (3).

**Lemma 2.3.** *The Chern character of $p_Y$ is given by $\tilde{\nu}^*\mathrm{ch}[p_T]$ and the Chern character of $p_T$ in cyclic homology is given by the formula*

$$\begin{aligned}\mathrm{ch}[p_T](z_0, z_1, \ldots, z_{2k}) &= (k!)^{-1}\mathrm{tr}_{\mathbb{C}^{2^n}}\left(\prod_{l=0}^{2k} p_T(z_l)\right) = \\ &= (k!)^{-1}\prod_{j=1}^{n}\prod_{l=1}^{2k}\frac{1+\bar{z}_{j,l}z_{j,l+1}}{1+|z_{j,l}|^2},\end{aligned}$$

*where we identify $z_{j,2k+1} = z_{j,0}$.*

*Proof.* We may write the Chern character of $p_T$ as the product of traces depending only on the $j$:th coordinate using (9). However, (8) implies that

$$\mathrm{tr}_{\mathbb{C}^2}\left(\prod_{l=0}^{2k} p_0(z_{j,l})\right) = \prod_{l=0}^{2k}\langle v(z_{j,l}), v(z_{j,l+1})\rangle = \prod_{l=0}^{2k}\frac{1+\bar{z}_{j,l}z_{j,l+1}}{1+|z_{j,l}|^2}.$$

□

# 3 Index theory for pseudo-differential operators

The index of an elliptic pseudo-differential operator can be expressed in terms of local formulas depending on the symbol via the Atiyah-Singer index theorem. In this section we will use elliptic pseudo-differential operators and the Atiyah-Singer index theorem to give an index formula for the degree of a continuous mapping. The theory of pseudo-differential operators can be found in [11] and for an introduction to the Atiyah-Singer index theorem we refer the reader to the survey article [1].

The Atiyah-Singer index theorem, see Theorem 1 of [1], states that the index of an elliptic pseudo-differential operator $A$ on the compact manifold $X$ without boundary can be calculated using de Rham cohomology as:

$$\mathrm{ind}\,(A) = \int_X \mathrm{ch}[A] \wedge Td(X),$$

where $\mathrm{ch}[A]$ is the Chern character of $A$ and $Td(X)$ is the Todd class of the complexified tangent bundle of $X$. Since the Chern character is a ring homomorphism we have the following lemma:

**Lemma 3.1.** *For a smooth projection valued function $p : X \to \mathcal{K}$ and a pseudo-differential operator $A$, the Chern character of the pseudo-differential operator $A_p := p(A \otimes id)p$ is given by $\mathrm{ch}[A_p] = \mathrm{ch}[A] \wedge \mathrm{ch}[p]$.*

Later on, in Theorem 4.5, the pseudo-differential operator will play a different role compared to the role in the Atiyah-Singer theorem. In the Atiyah-Singer theorem the elliptic pseudo-differential operator defines an element in $K$-theory which pairs with the $K$-homology class whose Connes-Chern character is the Todd class under the isomorphism (2) and gives an index. We will use the elliptic pseudo-differential operator in the dual way as a $K$-homology class that we pair with projections over $C^\infty(X)$ in terms of an index. The heuristic explanation of this method is that $K^0(X)$ is a ring and if $A$ is an elliptic pseudo-differential operator, $A$ defines both a $K$-homology class on $X$ and the symbol of $A$ defines an element $[A] \in K^0(X)$, using the Thom isomorphism, and $\mathrm{ind}_A(x) = \mathrm{ind}\,(x \cdot [A])$ for $x \in K^0(X)$ where $\mathrm{ind} : K^0(X) \to \mathbb{Z}$ is the index mapping of Atiyah-Singer. On the level of de Rham cohomology this is exactly the content of Lemma 3.1.

**Theorem 3.2.** *If $f : X \to Y$ is a smooth mapping between even-dimensional manifolds, $p_Y$ is as in* (11) *and $A$ is an elliptic pseudo-differential operator on $X$ we have the following degree formula:*

$$\mathrm{ch}_0[A]\deg(f) = \mathrm{ind}\,(A_{p_Y \circ f}) - \mathrm{ind}\,(A).$$

*If $A$ is of order 0, the same statement holds for continuous $f$.*

*Proof.* By Lemma 3.1, Theorem 2.2 and the Atiyah-Singer index theorem we have that

$$\begin{aligned}\mathrm{ind}\,(A_{p_Y \circ f}) &= \int_X \mathrm{ch}[f^* p_Y] \wedge \mathrm{ch}[A] \wedge Td(X) = \\ &= \int_X (1 + f^* \mathrm{d}V_Y) \wedge \mathrm{ch}[A] \wedge Td(X) = \\ &= \mathrm{ch}_0[A]\deg(f) + \int_X \mathrm{ch}[A] \wedge Td(X) = \\ &= \mathrm{ch}_0[A]\deg(f) + \mathrm{ind}\,(A).\end{aligned}$$

Since both $\mathrm{ind}\,(A_{p_Y \circ f})$ and $\deg(f)$ are well defined homotopy invariants for continuous $f$ when $A$ is of order 0 the final statement of the Theorem follows. □

To deal with analytic formulas for the mapping degree when $f$ is not smooth will require some more concrete information about Schatten class properties of pseudo-differential operators. We will restrict our attention to poly-homogeneous pseudo-differential operators.

**Lemma 3.3.** *A pseudo-differential operator $b$ of order $-1$ satisfies $b \in \mathcal{L}^q(L^2(X))$ for any $q > \dim(X)$.*

Lemma 3.3 is proved by using a rather standard technique for pseudo-differential operators. In Theorem 3.5, when we prove a similar result for Hölder continuous functions we will need some heavier machinery. We include a sketch of the proof of Lemma 3.3 just to highlight the difference in methods. Letting $\Delta_X$ denote the second order Laplace-Beltrami operator, the operator $(1-\Delta_X)^{1/2}b$ is of order 0 whenever $b$ is of order $-1$. Thus $b = (1-\Delta_X)^{-1/2}(1-\Delta_X)^{1/2}b$ and since $(1-\Delta_X)^{1/2}b$ is a bounded operator the Lemma follows if $(1-\Delta_X)^{-1/2}$ is in the Schatten class for any $q > \dim(X)$. This fact follows from the fact that the $k$:th eigenvalue of the Laplacian behaves like $k^{1/\dim(X)}$ as $k \to \infty$, a statement that goes back to [17].

**Lemma 3.4.** *The operator $b$ has an integral kernel $T \in C^\infty(X \times X \setminus D)$ satisfying the estimate $|T(x,y)| \lesssim |x-y|^{-\dim(X)}$ almost everywhere, if $b$ is a poly-homogeneous pseudo-differential operator of order 0 and $D$ denotes the diagonal in $X \times X$.*

Here we use the notation $a \lesssim b$ if there is a constant $C > 0$ such that $a \leq Cb$. Observe that the estimate on $T$ only holds for $x \neq y$ so the integral operator defined by $T$ must be realized as a principal value. We will not prove Lemma 3.4, but refer to Theorem 18.1.16 of [11] or Lemma 1.2.5 of [9].

**Theorem 3.5.** *If $F$ is a poly-homogeneous pseudo-differential operator of order 0 on a compact manifold $X$ without boundary and $a \in C^\alpha(X)$, then the operator $[F, \pi(a)]$ is Schatten class of order $q$ for any $q > \max(\dim(X)/\alpha, 2)$.*

We will prove Theorem 3.5 by two lemmas describing Schatten class properties of the commutator $[F, \pi(a)]$ in terms of its local nature as in Theorem 1.2. The first of the two lemmas is a direct consequence of Theorem 1.2:

**Lemma 3.6.** *Let $n/2 < \beta < n$ and suppose that $T$ is an integral operator on $\mathbb{R}^n$ whose integral kernel is bounded by $|x-y|^{-\beta}$, for any $\chi, \chi' \in C_c^\infty(\mathbb{R}^n)$ and $q > \max(n/(n-\beta), 2)$ the operator $\chi T \chi'$ is $q$-summable.*

*Proof.* First we observe that the condition $\beta < n$ implies that the kernel is locally integrable and the integral operator defined by $T$ can be defined without taking any principal values, so we may apply Russo's theorem directly to $\chi T \chi'$. Let us use the notation $K_\beta(x,y) := |x-y|^{-\beta}$. If $T$ has an integral kernel bounded by $K_\beta$ Theorem 1.2 implies that for some large $R > 0$ and for $q > 2$

$$\|\chi T \chi'\|_{\mathcal{L}^q(L^2(X))} \leq \left(\|\chi K_\beta \chi'\|_{q',q} \|\chi' K_\beta \chi\|_{q',q}\right)^{1/2} \lesssim$$
$$\lesssim \left( \int_{B(0,R)} \left( \int_{B(0,R)} \frac{\mathrm{d}V(x)}{|x-y|^{\beta q'}} \right)^{q/q'} \mathrm{d}V(y) \right)^{1/q} < \infty$$

if $\beta q' < n$ which is equivalent to $q > n/(n-\beta)$. □

**Lemma 3.7.** *Suppose that $F$ is a properly supported, poly-homogeneous pseudo-differential operator of order 0 in $\mathbb{R}^n$ and $a \in C^\alpha(\mathbb{R}^n)$ has compact support, then the operator $[F, \pi(a)]$ is Schatten class of order $q$ for any $q > \max(n/\alpha, 2)$.*

*Proof.* Since $F$ is a pseudo-differential operator of order 0, the operator $F$ can by Lemma 3.4 be represented by an integral kernel which is pointwise bounded by $|x-y|^{-n}$. Thus the integral kernel of $[F,\pi(a)]$ is bounded by $|a(x)-a(y)||x-y|^{-n}$. It follows from the Hölder continuity of $a$ that the integral kernel of $[F,\pi(a)]$ is pointwise bounded by the kernel $|x-y|^{-(n-\alpha)}$. While $F$ is properly supported and $a$ has compact support, we can take $\chi,\chi' \in C_c^\infty(\mathbb{R}^n)$ such that $\chi[F,\pi(a)]\chi' = [F,\pi(a)]$ and Lemma 3.6 implies that the operator $\chi[F,\pi(a)]\chi'$ is Schatten class of order $q$ for any $q > \max(n/\alpha, 2)$. □

Theorem 3.5 follows from Lemma 3.7 since Theorem 3.5 can be reduced to a local claim where Lemma 3.7 applies.

# 4 The index of a Hölder continuous twist

In this section we will combine Theorem 1.1 with Theorem 3.5 into index formulas for elliptic pseudo-differential operators twisted by Hölder continuous vector bundles. If $p : X \to \mathcal{K}(\mathbb{C}^N)$ is a continuous projection-valued function, we will use the notation $E_p$ for the vector bundle over $X$ corresponding to $p$ via Serre-Swan's theorem.

When $A$ is an elliptic pseudo-differential operator from the vector bundle $E$ to the vector bundle $E'$, we want to consider the, possibly unbounded, operator $A_p := p(A \otimes \mathrm{id}_{\mathbb{C}^N})p$ which is called the twist of $A$ by $p$ and acts between the vector bundle $E \otimes E_p$ and $E' \otimes E_p$. However, unless $A$ is of order 0, we must assume that $p$ is smooth to ensure that $A_p$ is a densely defined Fredholm operator. In the case $p$ is smooth, Lemma 3.1 and the Atiyah-Singer index theorem imply that

$$\mathrm{ind}\,(A_p) = \int_X \mathrm{ch}[p] \wedge \mathrm{ch}[A] \wedge Td(X),$$

and we also have the identity

$$\mathrm{ind}\,(A_p) = \mathrm{ind}\,(p(A(1+A^*A)^{-1/2} \otimes \mathrm{id}_{\mathbb{C}^N})p), \tag{14}$$

because $1 + A^*A$ is strictly positive. The right hand side of (14) is well defined for continuous $p$ and, as we will see, it can be calculated for Hölder continuous projections by means of Theorem 1.1 using a certain Fredholm module we associate with $A$. To construct this Fredholm module, we start by defining the odd, self-adjoint operator

$$\tilde{A} := \begin{pmatrix} 0 & A \\ A^* & 0 \end{pmatrix}$$

on $L^2(X, E \oplus E')$ which is graded by letting $L^2(X,E)$ be the even part and let $L^2(X,E)$ be the odd part. We define the mapping

$$\varphi : \mathbb{R} \to \mathbb{R}, \quad u \mapsto u(1+u^2)^{-1/2} \quad \text{and the operator} \quad F_A := \varphi(\tilde{A}).$$

The operator $F_A$ is an odd, self-adjoint operator of order 0. However, the square of the operator $F_A$ can be calculated as

$$F_A^2 = \tilde{A}^2(1+\tilde{A}^2)^{-1} = 1 - (1+\tilde{A}^2)^{-1} \neq 1.$$

To mend the problem $F_A^2 \neq 1$, we replace $\varphi$ by the function $\tilde{\varphi} : \mathbb{R} \to M_4(\mathbb{C})$ defined as

$$\tilde{\varphi}(u) := \begin{pmatrix} \varphi(u) & 0 & 0 & i(1+u^2)^{-1/2} \\ 0 & \varphi(u) & -i(1+u^2)^{-1/2} & 0 \\ 0 & i(1+u^2)^{-1/2} & -\varphi(u) & 0 \\ -i(1+u^2)^{-1/2} & 0 & 0 & -\varphi(u) \end{pmatrix}.$$

The function $\tilde{\varphi}$ satisfies $\tilde{\varphi}(u)^2 = 1$. If we equip $\mathbb{C}^4$ with the grading from the involution

$$\gamma_{\mathbb{C}^4} := 1 \oplus (-1) \oplus 1 \oplus (-1),$$

the operator $\tilde{F}_A := \tilde{\varphi}(\tilde{A})$ is an odd, self adjoint operator on the graded tensor-product $\mathbb{C}^4 \otimes L^2(X, E \oplus E')$. To simplify notations we set $\mathcal{E} := \mathbb{C}^4 \otimes (E \oplus E')$. The operator $\tilde{F}_A$ does satisfy that $\tilde{F}_A^2 = 1$. The operator $\tilde{F}_A$ can be written as a matrix of operators as

$$\tilde{F}_A := \begin{pmatrix} F_A & 0 & 0 & i(1+\tilde{A}^2)^{-1/2} \\ 0 & F_A & -i(1+\tilde{A}^2)^{-1/2} & 0 \\ 0 & i(1+\tilde{A}^2)^{-1/2} & -F_A & 0 \\ -i(1+\tilde{A}^2)^{-1/2} & 0 & 0 & -F_A \end{pmatrix}. \tag{15}$$

Since pseudo-differential operators are only pseudo-local, we will use a parameter $t$ to extract the singular part of $\tilde{F}_A$. Set $\tilde{F}(t) := \tilde{F}_{tA}$. Define $W_0$ as the smooth pseudo-differential operator given by the orthogonal finite-rank projection onto $\ker(\tilde{A})$ and set $W_0^\perp := 1 - W_0$. Observe that, since $\tilde{A}$ is elliptic, we have that $W_0 \in C^\infty(X, E \oplus E') \otimes_{alg} C^\infty(X, (E \oplus E')^*)$. We define

$$W := \begin{pmatrix} 0 & 0 & 0 & iW_0 \\ 0 & 0 & -iW_0 & 0 \\ 0 & iW_0 & 0 & 0 \\ -iW_0 & 0 & 0 & 0 \end{pmatrix}.$$

**Lemma 4.1.** *If $A$ is a poly-homogeneous, elliptic pseudo-differential operator of order at least $n/p$, the operator valued function $t \mapsto \tilde{F}(t)$ satisfies*

$$\|\tilde{F}(t) - \tilde{F} - W\|_{\mathcal{L}^p(L^2(X))} = \mathcal{O}(t^{-1}) \quad as \quad t \to \infty,$$

*where $\tilde{F}$ is the 0-homogeneous part of $\tilde{F}_A$.*

*Proof.* The operator $\tilde{F}(t) - \tilde{F} - W$ is a matrix consisting of terms of the form

$$\begin{aligned} a_1 &= i(1+t^2\tilde{A}^2)^{-1/2} - iW_0 = iW_0^\perp(1+t^2\tilde{A}^2)^{-1/2}W_0^\perp = \\ &= it^{-1}W_0^\perp\tilde{A}^{-1}W_0^\perp T_1 \qquad \text{and} \\ a_2 &= \tilde{A}(t(1+t^2\tilde{A}^2)^{-1/2} - W_0^\perp|\tilde{A}|^{-1}W_0^\perp) = \\ &= \tilde{A}(t(1+t^2\tilde{A}^2)^{-1/2} - |\tilde{A}|^{-1}) = t^{-1}W_0^\perp|\tilde{A}|^{-1}W_0^\perp T_2, \end{aligned}$$

for some $T_1, T_2 \in \mathcal{B}(L^2(X))$. Since both $a_1$ and $a_2$ are pseudo-differential operators of order lower than $-n/p$, it follows that their $\mathcal{L}^p$-norm behaves like $t^{-1}$ as $t \to \infty$ and the Lemma follows. □

We let $\pi : C^\alpha(X) \to \mathcal{B}(L^2(X, E \oplus E'))$ denote the even representation given by pointwise multiplication. Define $\tilde{\pi} : C^\alpha(X) \to \mathcal{B}(L^2(X, \mathcal{E}))$ by letting $\tilde{\pi} := \pi \oplus 0$ under the isomorphism $L^2(X, \mathcal{E}) \cong \mathbb{C}^4 \otimes L^2(X, E \oplus E')$.

**Theorem 4.2.** *If $A$ is a poly-homogeneous, elliptic operator of order at least $\alpha$ between two vector bundles $E$ and $E'$ over $X$ and $q > \max(\dim(X)/\alpha, 2)$, then the pair $(\tilde{\pi}, \tilde{F}_A)$ is a $q$-summable even Fredholm module over $C^\alpha(X)$.*

*Proof.* The operator $\tilde{F}_A$ is an elliptic, self-adjoint pseudo-differential operator of order 0 and $\tilde{F}_A^2 = 1$. Under the isomorphism $L^2(X, \mathcal{E}) \cong \mathbb{C}^4 \otimes L^2(X, E \oplus E')$ the decomposition (15) implies that

$$[\tilde{F}_A, \tilde{\pi}(a)] = \begin{pmatrix} [F_A, \pi(a)] & 0 & 0 & -i\pi(a)(1+\tilde{A}^2)^{-1/2} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -i(1+\tilde{A}^2)^{-1/2}\pi(a) & 0 & 0 & 0 \end{pmatrix}. \tag{16}$$

If the order of $A$ is bigger than $\alpha$, these facts together with Theorem 3.5 and Lemma 3.3 imply that $(\tilde{\pi}, \tilde{F}_A)$ is a $q$-summable even Fredholm module for any $q > \max(\dim(X)/\alpha, 2)$. □

Let us represent the pseudo-differential operator $\tilde{F}$ by the integral kernel $\tilde{K}_A$ which is a conormal distribution section of the big Hom-bundle $\mathrm{Hom}(\mathcal{E}, \mathcal{E})$. By (15), we can write $\tilde{K}_A = K_A \oplus K_A \oplus (-K_A) \oplus (-K_A)$ where $K_A$ is a conormal distribution section of the big Hom-bundle $\mathrm{Hom}(E \oplus E', E \oplus E')$. Observe that $K_A$ is defined by a smooth section $C^\infty(X \times X \setminus D, \mathrm{Hom}(E \oplus E', E \oplus E'))$ such that $|K_A(x, y)| \lesssim |x - y|^{-n}$.

We will use the notation $\Gamma_k$ for the subset of $\{1, 2, 3, 4\}^{2k}$ consisting of all sequences $(s_l)_{l=1}^{2k}$ satisfying the conditions that $s_1 \neq 4$, $s_{2k} \neq 3$ and $s_l = 3$ for some $l$ if and only if $s_{l+1} = 4$. These conditions are motivated by the form of the commutator (16). Let $w(I)$ denote the numeral of occurrences of 3 in $I$. Take $\Gamma_k^w \subseteq \Gamma_k$ as the subset of sequences with $w(I) = w$. To a sequence $I \in \Gamma_k$ we will associate the sequence $\Lambda(I) = (i_l)_{l=1}^{2k} \in \{1, 2, \ldots 2k\}^{2k}$ defined by

$$i_l = \begin{cases} l, & \text{if} \quad s_l = 1, 3 \\ l+1, & \text{if} \quad s_l = 2, 4 \end{cases},$$

where we identify $2k + 1$ with 1. For a sequence $I \in \Gamma_k$ we let $\iota(I) := (-1)^{\sum_{l=1}^{2k} i_l - l} \cdot i^{w(I)}$. With a projection-valued function $p : X \to \mathcal{K}$ and a sequence $I \in \Gamma_k$ we associate the function

$$Q_I^p(x_1, \ldots, x_{2k}) := \mathrm{tr}_{\mathbb{C}^{2^n}} \left( p(x_1) \prod_{l=1}^{2k} p(x_{i_l}) \right). \tag{17}$$

We also define

$$H_A(x_1, \ldots, x_{2k}) := \mathrm{str}_{E \oplus E'} \left( \prod_{l=1}^{2k} K_A(x_i, x_{i+1}) \right).$$

A straightforward calculation implies the following lemma:

**Lemma 4.3.** *For any projection-valued function $p : X \to \mathcal{K}(\mathbb{C}^N)$, the following identity holds:*

$$\mathrm{str}_{(E\oplus E')\otimes\mathbb{C}^N}\left(p(x_1)\prod_{j=1}^{2k}(p(x_{j+1})-p(x_j))K_A(x_j,x_{j+1})\right)= \\ =\sum_{(s_l)_{l=1}^{2k}\in\Gamma_k^0}\iota(I)Q_I^p(x_1,\ldots,x_{2k})H_A(x_1,\ldots,x_{2k}).$$

This Lemma describes the diagonal terms in the decomposition (16). To describe the products with the off-diagonal terms, we need some notation for the integral kernels. Set $K_1 = K_2 = K_A$ and $K_3 = K_4 = W_0$. For $I \in \Gamma_k$ we define the integral kernel

$$H_{A,I}(x_1,\ldots,x_{2k}) := \mathrm{str}_{E\oplus E'}\left(\prod_{l=1}^{2k}K_{s_l}(x_i,x_{i+1})\right).$$

**Lemma 4.4.** *The function*

$$(x_1,\ldots,x_{2k})\mapsto\sum_{(s_l)_{l=1}^{2k}\in\Gamma_k^w}\iota(I)Q_I^p(x_1,\ldots,x_{2k})H_{A,I}(x_1,\ldots,x_{2k}) \qquad (18)$$

*is absolutely integrable over $X^{2k}$ for all $w = 0, 1, \ldots, k$.*

This Lemma is a direct consequence of Theorem 3.5 and Theorem 1.3 since the sum over $\Gamma_k^w$ corresponds to the sum of the supertraces of the products between $p$ and $2(k-w)$ commutators between $p$ and $K_A$ and $2w$ commutators between $p$ and $W_0$. In fact, when $w > 0$ the integral of (18) will be the trace of a finite rank operator. One can decompose the function (18) even further by decomposing $\Gamma_k^w$ into equivalence classes under the equivalence relation $1 \sim 2$ and $3 \sim 4$, which again will be a decomposition into absolutely integrable functions.

**Theorem 4.5.** *Suppose that $A$ is an elliptic pseudo-differential operator of order at least $\alpha$ on the compact manifold $X$ without boundary, acting from $E$ to $E'$. If $p : X \to \mathcal{K}(\mathbb{C}^N)$ is a projection valued, Hölder continuous function of exponent $\alpha$ and $2k - 1 > \max(\dim(X)/\alpha, 2)$, the following index formula holds:*

$$\mathrm{ind}\,(p(F_{A,+}\otimes 1)p) = \langle \mathrm{cc}_k(\tilde{\pi},\tilde{F}_A), p\rangle_k = \\ =(-1)^k\int_{X^{2k}}\mathrm{str}\left(p(x_1)\prod_{j=1}^{2k}(p(x_{j+1})-p(x_j))K_A(x_j,x_{j+1})\right)\mathrm{d}V_{X^{2k}}+ \\ +(-1)^k\sum_{w=1}^{k}\int_{X^{2k}}\sum_{(s_l)_{l=1}^{2k}\in\Gamma_k^w}\iota(I)Q_I^p(x_1,\ldots,x_{2k})H_{A,I}(x_1,\ldots,x_{2k})\mathrm{d}V_{X^{2k}}$$

*where the trace is taken over $(E\oplus E')\otimes\mathbb{C}^N$ and we identify $x_1 = x_{2k+1}$. All of the integrals are absolutely convergent.*

Observe that if the dimension of $X$ is odd, $\mathrm{ind}\,(p(F_{A,+}\otimes 1)p) = 0$ for continuous $p$ since the Atiyah-Singer index theorem implies that it holds for smooth projection-valued functions.

*Proof.* The first equality follows from Theorem 1.1 and Theorem 4.2 since

$$\tilde{\pi}(p)(\tilde{F}_{A,+}\otimes 1)\tilde{\pi}(p) = (\pi(p)(F_{A,+}\otimes 1)\pi(p))\oplus 0.$$

Recall that, for any $t>0$, the operator $tA$ satisfies the same conditions as $A$ so by homotopy invariance of the Chern-Connes character and Lemma 4.1,

$$\begin{aligned}
&\langle \mathrm{cc}_k(\tilde{\pi},\tilde{F}_A),p\rangle_k = \langle \mathrm{cc}_k(\tilde{\pi},\tilde{F}_{tA}),p\rangle_k = \\
&= (-1)^k \mathrm{str}_{L^2(X,\mathcal{E}\otimes\mathbb{C}^N)}\left(\tilde{\pi}(p)[\tilde{F}_{tA},\tilde{\pi}(p)]\cdots[\tilde{F}_{tA},\tilde{\pi}(p)]\right) = \\
&= (-1)^k \mathrm{str}_{L^2(X,\mathcal{E}\otimes\mathbb{C}^N)}\left(\tilde{\pi}(p)[\tilde{F}+W,\tilde{\pi}(p)]\cdots[\tilde{F}+W,\tilde{\pi}(p)]\right) + \mathcal{O}(t^{-1}).
\end{aligned}$$

Since $p$ is Hölder continuous, $(p(x_{j+1})-p(x_j))K_A(x_j,x_{j+1})$ is locally integrable and of finite mixed $L^p$-norm by Theorem 3.5. Let us set $\tilde{p}:=\tilde{\pi}(p)$ and $\tilde{K}_A^W := \tilde{K}_A + W$. Theorem 1.2 and the calculations above imply the equality

$$\begin{aligned}
&\mathrm{str}_{L^2(X,\mathcal{E}\otimes\mathbb{C}^N)}\left(\tilde{\pi}(p)[\tilde{F}+W,\tilde{\pi}(p)]\cdots[\tilde{F}+W,\tilde{\pi}(p)]\right) = \\
&= \int_{X^{2k}} \mathrm{str}\left(\tilde{p}(x_1)\prod_{j=1}^{2k}\left(\tilde{p}(x_{j+1})\tilde{K}_A^W(x_j,x_{j+1}) - \tilde{K}_A^W(x_j,x_{j+1})\tilde{p}(x_j)\right)\right)\mathrm{d}V_{X^{2k}} \\
&= \int_{X^{2k}} \sum_{(s_l)_{l=1}^{2k}\in\Gamma_k} \iota(I)Q_I^p(x_1,\ldots,x_{2k})H_{A,I}(x_1,\ldots,x_{2k})\mathrm{d}V_{X^{2k}}.
\end{aligned}$$

While the term $\langle \mathrm{cc}_k(\tilde{\pi},\tilde{F}_A),p\rangle_k$ is constant, the second identity stated in the theorem follows from Lemma 4.3 by letting $t\to\infty$. □

# 5 Degrees of Hölder continuous mappings

Returning now to the degree calculations, we will use Theorem 4.5 for a particular choice of pseudo-differential operator. Assume that $X$ is a compact Riemannian manifold of dimension $2n$ without boundary. The Hodge grading on $\bigwedge^* T^*X$ is defined by the involution $\tau$ defined on a $p$-form $\omega$ by

$$\tau\omega = i^{p(p-1)+n} * \omega,$$

where $*$ denotes the Hodge duality. Observe that if $(e_j)_{j=1}^{2n}$ is an oriented, orthonormal basis of the cotangent space, the operator $\tau$ can be written as

$$\tau = \left(\frac{i}{2}\right)^n \prod_{j=1}^{2n}(e_j\wedge -e_j\neg). \tag{19}$$

We let $E_+$ denote the sub-bundle of $\bigwedge^* T^*X$ consisting of even vectors with respect to the Hodge grading and $E_-$ the sub-bundle of odd vectors with respect to the Hodge grading. The operator $\tau$ anti-commutes with $\mathrm{d}+\mathrm{d}^*$ so $A=\mathrm{d}+\mathrm{d}^*$ is a well defined operator from $E_+$ to $E_-$. The operator $A$ is called the signature operator. Observe that $\tilde{A}=\mathrm{d}+\mathrm{d}^*$ as an operator on $\bigwedge^* T^*X$ and $\tilde{A}^2$ is the Laplace-Beltrami operator on $X$. By Theorem 4.2 the pair $(\tilde{\pi},\tilde{F}_{\mathrm{d}+\mathrm{d}^*})$ is a $q$-summable Fredholm module over $C^\alpha(X)$.

Assume that $f : X \to Y$ is a Hölder continuous function where $Y$ is a $2n$-dimensional manifold. We can choose an open subset $U \subseteq Y$ such that there is a diffeomorphism $\nu : U \to B_{2n}$. In fact, by the closed mapping lemma, since $X$ is compact and $Y$ Hausdorff, the mapping $f$ is open and we can take $U$ in such a way that there is an open set $U_0 \subseteq X$ satisfying $U \subseteq f(U_0)$ and $E_+$ and $E_-$ are trivial over $U_0$.

Taking $\tilde{\nu} : Y \to \mathbb{C}^n$ as the Lipschitz continuous function defined in (12) we can for $k > n/\alpha$ define the integrable function $\tilde{f}_k \in L^1(X^{2k})$ as:

$$\tilde{f}_k(x_1, \dots, x_{2k}) := \tag{20}$$

$$= \sum_{I \in \Gamma_k} \iota(I) Q_I^{p_T}(\tilde{\nu} f(x_1), \dots, \tilde{\nu} f(x_{2k})) H_{\mathrm{d}+\mathrm{d}^*,I}(x_1, \dots, x_{2k}) = \tag{21}$$

$$= \sum_{w=0}^{k} \sum_{I \in \Gamma_k^w} \iota(I) H_{\mathrm{d}+\mathrm{d}^*,I}(x_1, \dots, x_{2k}) \prod_{j=1}^{n} \prod_{l=0}^{2k} \frac{1 + \overline{f(x_{i_l})}_j f(x_{i_{l+1}})_j}{1 + |f(x_{i_l})_j|^2}, \tag{22}$$

where the second formula follows from Lemma 2.3. The kernel $H_{\mathrm{d}+\mathrm{d}^*,I}$ is in general quite hard to find. In local coordinates, the operator $K_{\mathrm{d}+\mathrm{d}^*}$ will be similar to a Riesz transform. The operator $W_0$ is the projection onto the finite-dimensional space of harmonic forms on $X$. We will demonstrate this by calculating the kernels $H_{\mathrm{d}+\mathrm{d}^*,I}$ explicitly on $S^{2n}$ in the next section.

**Theorem 5.1.** *Suppose that $X$ and $Y$ are smooth compact connected manifolds without boundary of dimension $2n$ and $f : X \to Y$ is Hölder continuous of exponent $\alpha$. When $k > n/\alpha$ the following integral formula holds:*

$$\deg(f) = -\mathrm{sign}(X) + (-1)^k \int_{X^{2k}} \tilde{f}_k(x_1, \dots, x_{2k}) \mathrm{d}V_{X^{2k}}$$

*where $\tilde{f}_k$ is as in* (20)–(22).

*Proof.* While $F_{\mathrm{d}+\mathrm{d}^*} = (\mathrm{d} + \mathrm{d}^*)(1 + \Delta)^{-1/2}$, we have that

$$\mathrm{ind}\,(F_{\mathrm{d}+\mathrm{d}^*,+}) = \mathrm{ind}\,(\mathrm{d} + \mathrm{d}^*) = \mathrm{sign}(X).$$

The operator $\mathrm{d} + \mathrm{d}^*$ satisfies $\mathrm{ch}_0[\mathrm{d} + \mathrm{d}^*] = \hat{L}_0(T^*X) = 1$, the constant term in the $\hat{L}$-genus. Therefore Theorem 3.2 and Theorem 4.5 implies that

$$\begin{aligned}
\deg(f) &= \mathrm{ind}\,((p_Y \circ f \otimes \mathrm{id}) F_{\mathrm{d}+\mathrm{d}^*,+}(p_Y \circ f \otimes \mathrm{id})) - \mathrm{ind}\,(F_{\mathrm{d}+\mathrm{d}^*,+}) = \\
&= \langle \mathrm{cc}_k(\tilde{\pi}, \tilde{F}_A), f^* p_Y \rangle_k - \mathrm{sign}(X) = \\
&= (-1)^k \int_{X^{2k}} \tilde{f}_k(x_1, \dots, x_{2k}) \mathrm{d}V_{X^{2k}} - \mathrm{sign}(X).
\end{aligned}$$

□

A couple of remarks on the choice of $A$ as the signature operator are in order. This choice is rather superfluous since any pseudo-differential operator $A$ of order 1 with $\mathrm{ch}_0[A] \neq 0$ will give a degree formula similar to that in Theorem 5.1. If one can find an invertible $A$ on $X$ such that $\mathrm{ch}_0[A] \neq 0$, the formula of Theorem 5.1 would be much simpler since $\tilde{f}_k$ will not contain any contributions from $\Gamma_k \setminus \Gamma_k^0$.

The signature operator has been studied on Lipschitz manifolds, see [15], which gives an analytic degree formula for Hölder continuous mappings between even-dimensional Lipschitz manifolds. On Lipschitz manifolds, the Atiyah-Singer theorem is replaced by Teleman's index theorem from [16]. Of course, there are some analytic difficulties in the proof of Theorem 3.5 for Lipschitz manifolds, that more or less manifests themselves on a notational level.

# 6 Example on $S^{2n}$

Let us end this paper by writing down the integral kernels in the case of a Hölder continuous function $f : S^{2n} \to Y$ where $Y$ is a compact, connected $2n$-dimensional manifold without boundary. We will compare the method of using pseudo-differential operators from Theorem 5.1 with that of using Henkin-Ramirez kernels from [8]. The degree formula of [8] is based on the usage of the cocycle (5).

To apply Theorem 5.1 the kernels $H_{\mathrm{d}+\mathrm{d}^*,I}$ need to be calculated. Define $U_0 : L^2(S^{2n}, \bigwedge T^*S^{2n}) \to L^2(\mathbb{R}^{2n}, \bigwedge T^*\mathbb{R}^{2n})$ by pulling back along the mapping $\lambda : \mathbb{R}^{2n} \to S^{2n}$ defined by

$$\mathbb{R}^{2n} \ni x \mapsto \left( \frac{|x|^2 - 1}{|x|^2 + 1}, \frac{2x}{|x|^2 + 1} \right) \in S^{2n} \subseteq \mathbb{R}^{2n+1}$$

and equipping $\mathbb{R}^{2n}$ with the pull-back metric. Since the metric is positive definite we can define the unitary mapping $U : L^2(S^{2n}, \bigwedge T^*S^{2n}) \to L^2(\mathbb{R}^{2n}) \otimes \mathbb{C}^{2^{2n}}$ by composing $U_0$ with the unitary mapping $L^2(\mathbb{R}^{2n}, \bigwedge T^*\mathbb{R}^{2n}) \to L^2(\mathbb{R}^{2n}) \otimes \mathbb{C}^{2^{2n}}$ defined by the metric. Clearly, we have that $UK_{\mathrm{d}+\mathrm{d}^*}U^*$ is a pseudo-differential operator on $\mathbb{R}^{2n}$ with symbol $\xi \mapsto i(\xi \wedge -\xi\neg)/\sqrt{2}|\xi|$, if we identify $\mathbb{C}^{2^{2n}}$ with $\bigwedge \mathbb{C}^{2n}$. For any $a \in C^\alpha(S^{2n})$ we have that $UaU^* = a \circ \lambda$.

Let us find the integral kernel $K_1$ of $UK_{\mathrm{d}+\mathrm{d}^*}U^*$. Since the symbol of $K_1$ is a homogeneous function that commutes with the $SU(2n)$-action on $\bigwedge \mathbb{C}^{2n}$, the integral kernel $K_1$ is given by

$$K_1(x, y) = c_n \frac{(x - y) \wedge -(x - y)\neg}{\sqrt{2}|x - y|^{2n+1}}$$

for some constant $c_n$. So the operator defined by $K_1$ is a matrix of Riesz transforms. While the harmonic forms on $S^{2n}$ are spanned by the constant function and the volume form, the kernel $W_0$ is the constant projection $\bigwedge T^*S^{2n} \to \mathbb{C} \oplus \bigwedge^{2n} T^*S^{2n}$. So we can write the kernel $K_3$ of $UW_0U^*$ as

$$K_3(x, y) = g(x)g(y)(1 + \prod_{j=1}^{2n} e_j \wedge e_j\neg),$$

where $g(x) := c_n'(1 + |x|^2)^{-n}$ for some constant $c_n'$. Using these expressions for $K_1$ and $K_3$ it follows from (19) that

$$H_{\mathrm{d}+\mathrm{d}^*,I}(x_1, \ldots, x_{2k}) = \mathrm{str}\left( \prod_{l=1}^{2k} K_{s_l}(x_l, x_{l+1}) \right) = \mathrm{tr}\left( \tau \prod_{l=1}^{2k} K_{s_l}(x_l, x_{l+1}) \right) =$$

$$= \sum_{\sigma \in S_{2(k-w)}} \sum_{m=0}^{n} i^n \mathrm{sign}(\sigma) c_n^{k_I} H_{\sigma,m,I}(x_1, \ldots, x_{2k}),$$

when $w(I) = w$ and where

$$H_{\sigma,m,I}(x_1,\ldots,x_{2k}) := \prod_{s_l\neq 1,2} g(x_l)g(x_{l+1}) \prod_{l=1,s_l=1,2}^{m} \frac{\langle e_j, x_{\sigma(l)} - x_{\sigma(l)+1}\rangle}{|x_{\sigma(l)} - x_{\sigma(l)+1}|^{2n+1}} \cdot$$
$$\cdot \prod_{l=m+1,s_l=1,2}^{k} \frac{\langle x_{\sigma(l)} - x_{\sigma(l)+1}, x_{\sigma(l+1)} - x_{\sigma(l+1)+1}\rangle}{|x_{\sigma(l)} - x_{\sigma(l)+1}|^{2n+1}|x_{\sigma(l+1)} - x_{\sigma(l+1)+1}|^{2n+1}}.$$

By this notation we mean that an element $\sigma$ in the symmetric group $S_{2(k-w)}$, on $2(k-w)$ elements, acts on the indices $l$ such that $s_l = 1, 2$. Here we use the notation $\langle\cdot,\cdot\rangle$ for the scalar product. For a Hölder continuous function $f : S^{2n} \to Y$ we take an open set $U \subseteq Y$ such that there is a diffeomorphism $\nu : U \to B_{2n}$ and consider the Hölder continuous function $f_0 := \tilde{\nu} f \lambda : \mathbb{R}^{2n} \to S^{2n}$, where $\tilde{\nu}$ is as in equation (12). The signature of a sphere is 0 so Theorem 5.1 and (22) implies the following degree formula:

$\deg(f) =$

$$= (-1)^k \sum_{w=0}^{k} \int_{\mathbb{R}^{4nk}} \sum_{I\in\Gamma_k^w} \sum_{\sigma\in S_{2(k-w)}} \sum_{m=0}^{n} i^n \mathrm{sign}(\sigma) c_n^{k_I} \iota(I) \cdot$$
$$\cdot \prod_{j=1}^{n}\prod_{l=1}^{2k} \frac{1 + \overline{f_0(x_j)}_{i_l} f_0(x_j)_{i_{l+1}}}{1 + |f_0(x_j)_{i_l}|^2} \cdot$$
$$\cdot \prod_{s_l\neq 1,2} g(x_l)g(x_{l+1}) \prod_{l=1,s_l=1,2}^{m} \frac{\langle e_j, x_{\sigma(l)} - x_{\sigma(l)+1}\rangle}{|x_{\sigma(l)} - x_{\sigma(l)+1}|^{2n+1}} \cdot$$
$$\cdot \prod_{l=m+1,s_l=1,2}^{k} \frac{\langle x_{\sigma(l)} - x_{\sigma(l)+1}, x_{\sigma(l+1)} - x_{\sigma(l+1)+1}\rangle}{|x_{\sigma(l)} - x_{\sigma(l)+1}|^{2n+1}|x_{\sigma(l+1)} - x_{\sigma(l+1)+1}|^{2n+1}} \mathrm{d}V_{\mathbb{R}^{2nk}},$$

where $f_0(x)_i$ denotes the $i$:th complex coordinate of $f_0(x)$ if $f_0(x) \in U$ and $\infty$ if $f_0(x) \notin U$.

If we attempt using the degree formula of [8] to a function $f : S^{2n} \to Y$ we must in some way change the dimension. We will do so by finding a strictly pseudo-convex domain $\Omega$ in $\mathbb{C}^{n+1}$ such that $\partial\Omega = S^{2n} \times S^1$ and use the degree formula of [8] to calculate the degree of $f \times \mathrm{id} : S^{2n} \times S^1 \to Y \times S^1$. To find such domain $\Omega$ we define the function $\rho \in C^\infty(\mathbb{C}^{n+1})$ as

$$\rho(z_1,\ldots,z_{n+1}) := 4|1 - z_1 z_2|^2 + |z|^2 - 3.$$

We let $\Omega := \{z \in \mathbb{C}^{n+1} : \rho(z) < 0\}$. Clearly, $\rho$ is strictly pluri-subharmonic in $\Omega$ and $\mathrm{d}\rho \neq 0$ on $\partial\Omega$. Furthermore

$$\rho(z) = 4(1 - 2\Re(z_1 z_2) + |z_1 z_2|^2) + |z|^2 - 3.$$

So $\Omega$ is a relatively compact strictly pseudo-convex domain in $\mathbb{C}^{n+1}$ with smooth boundary. Writing down the function $\rho$ in its real argument verifies that $\Omega \cong B_{2n+1} \times S^1$ and $\partial\Omega \cong S^{2n} \times S^1$.

Let us find the Henkin-Ramirez kernel $H_R$ of the boundary $\partial\Omega$. In order to do this we will use the Fornaess embedding theorem, see [7]. This approach to construct the Henkin-Ramirez kernel goes as follows; if we find a proper, holomorphic mapping from $\Omega$ into a convex domain, we obtain the holomorphic support function of $\partial\Omega$ by pulling back the support function in the convex domain. Define the holomorphic function $\Psi : \mathbb{C}^{n+1} \to \mathbb{C}^{n+2}$ by $(z_1, z_2, \ldots, z_{n+1}) \mapsto (2(1 - z_1 z_2), z_1, z_2, \ldots, z_{n+1})$. Clearly $\Omega$ is mapped in to the ball of radius 3, $3B_{2n+4}$, and $\Omega = \Psi^{-1}(3B_{2n+4})$. Therefore $\Omega$ has the support function

$$a(z,\zeta) := (\Psi(\zeta) - \Psi(z)) \cdot \overline{\Psi(\zeta)} = \bar{\zeta} \cdot (\zeta - z) + 4(\zeta_1\zeta_2 - z_1 z_2)\overline{\zeta_1\zeta_2}. \tag{23}$$

Thus the integral kernel of the Henkin-Ramirez kernel for $S^{2n} \times S^1$ is given by

$$H_R(z,\zeta)\mathrm{d}V_{S^{2n}\times S^1} = \frac{1}{(2\pi i)^n} \frac{s \wedge (\bar{\partial}_\zeta s)^{2(n-1)}}{a(z,\zeta)^n},$$

where

$$s(z,\zeta) = \partial|\zeta|^2 + 2(1 - \overline{\zeta_1\zeta_2})(z_2\mathrm{d}\zeta_1 + \zeta_1\mathrm{d}\zeta_2).$$

Because of (23) the function $a(z,\zeta)$ is symmetric in the sense that for $z, \zeta \in \partial\Omega$ we have $a(z,\zeta) = \overline{a(\zeta,z)}$. However, the kernel $H_R$ is not symmetric as is seen from the expression for $s$.

Concluding the case of a Hölder continuous function $f : S^{2n} \to Y$, we take an open subset $U \subseteq Y$ diffeomorphic to a ball and choose a diffemorphism $\nu : U \times (S^1 \setminus \{pt\}) \to B_{2n+1}$. If we let $f_1 : S^n \times S^1 \to S^{n+1}$ be the Hölder continuous function constructed by extending $\nu(f \times \mathrm{id})$ as in (12), Theorem 4.4 of [8] implies the degree formula

$$\deg(f) = \frac{(-1)^n}{(2\pi i)^{2kn}} \int_{(S^{2n}\times S^1)^{2k+1}} \sum_{\sigma \in S_{2(2k-1)}} \sum_{l=0}^{2k-1} c_{l,\sigma} \prod_{i=1}^{l} \langle f_1(z_{\sigma(2i-1)}), f_1(z_{\sigma(2i)})\rangle \cdot \\ \cdot \prod_{j=0}^{2k} \frac{s(z_j, z_{j+1}) \wedge (\bar{\partial}_{z_{j+1}} s(z_j, z_{j+1}))^{2(n-1)}}{a(z_j, z_{j+1})^n},$$

where $c_{l,\sigma} = (-1)^l 2^{n-l-1}\varepsilon_l(\sigma)$ and $\varepsilon_l(\sigma)$ is the order parity of $\sigma$.

# References


[1] M. F. Atiyah, I.M. Singer, *The index of elliptic operators on compact manifolds*, Bull. Amer. Math. Soc. 69 1963 422–433.

[2] B. Blackadar, *K-theory for operator algebras*, Second edition. Mathematical Sciences Research Institute Publications, 5. Cambridge University Press, Cambridge, 1998.

[3] A. Benedek, R. Panzone, *The spaces $L^P$, with mixed norm*, Duke Math. J. Volume 28, Number 3 (1961), 301-324.

[4] A. Connes, *Noncommutative differential geometry*, Inst. Hautes Etudes Sci. Publ. Math. No. 62 (1985), 257–360.

[5] A. Connes, *Noncommutative geometry*, Academic Press, Inc., San Diego, CA, 1994.

[6] J. Cuntz, R. Meyer, J.M. Rosenberg, *Topological and Bivariant K-theory*, Birkhäuser 2007.

[7] J.E. Fornaess, *Embedding strictly pseudoconvex domains in convex domains*, Amer. J. Math. 98 (1976), no. 2, 529–569.

[8] M. Goffeng, *Analytic formulas for topological degree of non-smooth mappings: the odd-dimensional case*, arXiv:1004.1018v1.

[9] P.B. Gilkey, *Invariance theory, the heat equation, and the Atiyah-Singer index theorem*, Mathematics Lecture Series, 11. Publish or Perish, Inc., Wilmington, DE, 1984.

[10] E. Guentner, N. Higson, *A note on Toeplitz operators*, Internat. J. Math. 7 (1996), no. 4, 501–513.

[11] L. Hörmander, *The analysis of linear partial differential operators. III. Pseudo-differential operators*, Reprint of the 1994 edition. Classics in Mathematics. Springer, Berlin, 2007.

[12] D. O'Regan, Y.J. Cho, Y-Q. Chen, *Topological degree theory and applications*, Series in Mathematical Analysis and Applications, 10. Chapman & Hall/CRC, Boca Raton, FL, 2006

[13] B. Russo, *On the Hausdorff-Young Theorem for Integral Operators*, Pacific Journal of Mathematics, Vol 68, No. 1, 197.

[14] R.G. Swan, *Vector Bundles and Projective Modules*, Transactions of the American Mathematical Society 105 (2): 264-277.

[15] N. Teleman, *The index of signature operators on Lipschitz manifolds*, Inst. Hautes Etudes Sci. Publ. Math. No. 58 (1983), 39–78 (1984).

[16] N. Teleman, *The index theorem for topological manifolds*, Acta Math. 153 (1984), no. 1-2, 117–152.

[17] H. Weyl, *Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen (mit einer Anwendung auf die Theorie der Hohlraumstrahlung)*, Math. Ann. 71 (1912), no. 4, 441–479.